

Undiscovered Petroleum Resources for the Woodford Shale and Thirteen Finger Limestone–Atoka Shale Assessment Units, Anadarko Basin—Continued




Figure 17. Oil production from the Thirteen Finger limestone is partly within the start of oil generation based on a transformation ratio (TR) of 0.1%. The AU boundary approximates the extent of maturation for gas generation based on the 1.2% R_o and 99% TR contours. Perryton and Knowles NW field information is from IHS Energy (2009a, 2009b).




Figure 18. Isopach from the top of the Thirteen Finger limestone to the Morrow Group. Red lines are figure 19 cross section locations. White hexagons are Perryton (Texas) and Knowles (Oklahoma panhandle) oil fields. Morrow and Thirteen Finger limestone tops are based on Andrews (1999), examination of more than 120 well logs, and edited tops from IHS (2009 a, b). Morrow Formation faults (green) are modified from Andrews (1999).





Figure 19. Southwest to northeast (A-A') and southeast to northwest (B-B') cross sections of the Thirteen Finger limestone (TRFG) of the Pennsylvanian Atoka Group (ATOKA). Lines of section are on figure 18. Vertical scale is feet relative to the base of the Thirteen Finger limestone. MRMT is Marmaton Group, CHRK is Cherokee Group, MRRW is Morrow Group, and MSSP is top of the Mississippian.

Conclusions

Paleozoic source rocks are thermally mature for oil and gas generation across most of the Anadarko Basin in Oklahoma and Texas. The deep basin of Oklahoma and Texas is thermally mature for gas from Atokan and older source rocks. Deep basin areas that are overmature for gas generation are within the gas preservation window, as indicated by gas production from Woodford Shale and older formations in this area, and on modeled levels of thermal maturation. Almost all of the Colorado and Kansas portions of the basin are thermally immature for petroleum generation. Source of oil and gas in these areas is long-distance migration radially northward from the deep basin in Oklahoma and proximal Texas, and northward from the panhandle area of Texas.

Acknowledgments

Michael Lewan provided the comparison chart of Woodford Shale hydrous pyrolysis transformation ratios and vitrinite reflectance; I also have benefited greatly from his knowledge and research. Chesapeake Energy has been very generous with their time, data, and expertise, which is gratefully acknowledged. The author would like to thank Kathy Winner of IHS Energy and IHS Energy for providing information for our mapping and modeling efforts.

References

- Adler, F.J., Caplan, W.M., Carlson, M.P., Goebel, E.D., Henslee H.T., Hick, I.C., Larson, T.G., McCracken, M.H., Parker, M.C., Rascoe, G., Jr., Schramm, M.W., and Wells, J.S., 1971, Future petroleum provinces of the Mid-Continent, in Cram, I.H., ed., Future petroleum provinces of the United States—their geology and potential: American Association of Petroleum Geologist Memoir 15, v. 2, p. 985–1120.
- Andrews, R.D., 1999, Map showing regional structure at the top of the Morrow Formation in the Anadarko Basin and shelf of Oklahoma: Oklahoma Geologic Survey OGS SP 99 pl. 4.
- Burruss, R.C., and Hatch, J.R., 1989, Geochemistry of oils and hydrocarbon source rocks, greater Anadarko Basin—Evidence for multiple sources of oils and long-distance oil migration, in Johnson, K.S., ed., Anadarko Basin symposium, 1988: Oklahoma Geological Survey Circular 90, p. 53–54.
- Cardott, B.J., 1989, Thermal maturation of the Woodford Shale in the Anadarko Basin, in Johnson, K.S., ed., Anadarko Basin symposium, 1988: Oklahoma Geological Survey Circular 90, p. 32–46.
- Gallardo, Jaquidon, and Blackwell, D.D., 1997, Thermal structure of the Anadarko Basin: American Association of Petroleum Geologists Bulletin, v. 83, no. 2, p. 333–361.
- Higley, D.K., Gaswirth, S.B., Abbott, M.M., Charpentier, R.R., Cook, T.A., Ellis, G.S., Gianoutsos, N.J., Hatch, J.R., Klett, T.R., Nelson, Philip, Pawlewicz, M.J., Pearson, O.N., Pollastro, R.M., and Schenk, C.J., 2011, Assessment of undiscovered oil and gas resources of the Anadarko Basin Province of Oklahoma, Kansas, Texas, and Colorado, 2010: U.S. Geological Survey Fact Sheet 2011-3003, 2 p. accessed on Feb. 2011, at <http://pubs.usgs.gov/fs/2011/3003/>.
- Howery, S.D., 1993, A regional look at Hunton production in the Anadarko Basin, in Johnson, K.S., ed., Hunton Group core workshop and field trip: Oklahoma Geological Survey Special Publication 93–4, p. 77–81.
- IHS Energy, 2009a, IHS Energy Well database: Database available from IHS Energy, 15 Inverness Way East, Englewood, CO 80112.
- IHS Energy, 2009b, GDS database: Geological Data Services database available from IHS Energy, 15 Inverness Way East, Englewood, CO 80112.
- Lewan, M.D., and Ruble, T.E., 2002, Comparison of petroleum generation kinetics by isothermal hydrous and nonisothermal open-system pyrolysis: Organic Geochemistry, v. 33, no. 12, p. 1457–1475.
- Price, L.C., 1997, Minimum thermal stability levels and controlling parameters of methane, as determined by C15+ hydrocarbon thermal stabilities, in Dyman, T.S., Rice, D.D., and Westcott, P.A., eds., Geologic controls of deep natural gas resources in the United States: U.S. Geological Survey Bulletin 2146-K, p. 135–176.
- Rottmann, Kurt, 2000a, Structure map of Hunton Group in Oklahoma and Texas Panhandle: Oklahoma Geologic Survey SP 2000–2, pl. 3.
- Rottmann, Kurt, 2000b, Isopach map of Woodford Shale in Oklahoma and Texas Panhandle: Oklahoma Geologic Survey SP 2000–2, pl. 2.
- Schlumberger Ltd., 2011, PetroMod Basin and Petroleum Systems Modeling Software: IES GmbH, Ritterstrasser, 23, 52072 Aachen, Germany, accessed on Jan. 2011, at <http://www.ies.de>
- Sweeney, J.J., and Burnham, A.K., 1990, Evaluation of a simple model of vitrinite reflectance based on chemical kinetics: American Association of Petroleum Geologists Bulletin, v. 74, p. 1559–1570.